


## Factorization for DIS

To finish the lecture, let us briefly discuss factorization for DIS for large  $M_x^2$ , which is slightly simpler than the Surlakov form factor discussed above.

Pictorially, it looks as follows:



hadron final state

$$P_x = (p + q)^2$$

Similarly to our treatment of  $\sigma(e^+e^- \rightarrow X)$  one can obtain the DIS cross section from the imaginary part of the expectation value of two vector currents


$$T_{\mu\nu}(p, q) = \frac{1}{2} \sum_{\text{spin}} i \int d^4x e^{iqx} \overbrace{\langle P(p) | T \{ J_\mu(x) J_\nu(0) \} | P(p) \rangle}^{\text{average over proton spin}}$$

$$\langle P(p) | T \{ J_\mu(x) J_\nu(0) \} | P(p) \rangle$$

Proton  $\nearrow$

$$J^\mu = \sum_q e_q \bar{\psi}_q \gamma^\mu \psi_q$$

Pictorially:



$L^{\mu\nu}$

$x$

$\text{Im}[T_{\mu\nu}]$

As for the R-ratio, we then want to match

$$T \{ \mathcal{J}_\mu(x) \mathcal{J}_\nu(0) \} = \sum_i C_i O_i$$

↑  
Operators in SCET

To write down the operators, we choose the Breit frame, in which

$$q^\mu = (0, 0, 0, -Q) = \frac{Q}{2} (\bar{n}^\mu - n^\mu)$$

$$p^\mu = \overset{O(2)}{\bar{n} \cdot p} \frac{\bar{n}^\mu}{2} + \overset{O(\lambda^2)}{\frac{\mu_p^2}{\bar{n} \cdot p}} \frac{n^\mu}{2}$$

↑  
large,  $O(Q)$

↑  $p^2 = \mu_p^2 !$

The proton is described by a collinear field along the  $w$ -direction and we should write down all operators which have a nontrivial spin-averaged proton matrix element. The leading power operators are of the form

$$O_q = \bar{\chi}_c(s\bar{n}) \Gamma \chi_c(t\bar{n})$$

$$\Gamma = \frac{\not{1}}{2}, \frac{\not{1}}{2} \gamma_5, \frac{\not{1}}{2} \gamma_\perp^m$$

↑  
only this one

↖ Dirac basis between  $\chi_c$  fields.

gives nonzero matrix element

and (note  $\bar{n} \cdot A = 0$ )

$$O_g = A_c^\mu(s\bar{n}) A_c^\nu(t\bar{n}) \Gamma_{\mu\nu}$$

$$\Gamma_{\mu\nu} \in \{g_{\mu\nu}^\perp, n_\mu n_\nu\}$$

↑  
only this  
appears.

↑  
power suppressed

The product of currents thus has  
the form

$$T \{ J_\mu(z) J_\nu(0) \}$$

collinear  
quark field  
for flavor  $q$   
↓

can shift one  
field to zero using  
transl. invariance  
↓

$$= \sum_q \int ds C_{\mu\nu}^q(z, s) \bar{\chi}_q(s\bar{n}) \frac{1}{2} \chi_q(0)$$

$$- \int ds C_{\mu\nu}^g(z, s) A_{1\mu}^{\alpha, a}(s\bar{n}) A_{1\nu}^a(0)$$

$$+ O(\lambda^3)$$

This is reminiscent of the OPE, except that the operators are smeared along the  $n$  lightcone, which reflects the fact that the  $\bar{n} \cdot \partial$  derivatives are unsuppressed.

Note that the soft fields decouple also from the operators  $O_q$  and  $O_g$ . To see this note that

$$\begin{aligned} \chi_c(x) &\rightarrow S_n(x_-) \chi_c^{(0)}(x) \\ &\rightarrow \chi_c(s\bar{n}) \rightarrow S_n(0) \chi_c(s\bar{u}) \end{aligned}$$

$\bar{n} \cdot x \frac{u^+}{2}$   
 $\downarrow$

Therefore only collinear fields contribute.

The matrix elements of  $O_q$  &  $O_{\bar{q}}$  define the Parton Distribution Functions

$$\begin{aligned} & \frac{1}{2} \sum_{\text{spin}} \langle P(p) | \bar{\chi}_q(s\bar{u}) \not{\epsilon} \chi_q(0) | P(p) \rangle \\ &= \bar{n} \cdot p \int_{-1}^1 d\xi f_{q/p}(\xi) e^{i\xi s\bar{u} \cdot p} \end{aligned}$$

⌈ Note:  $f_{\bar{q}/p}(\xi) = -f_{q/p}(-\xi)$

is the anti-quark PDF

⌋

Intuitively, the PDF can be understood by noting that  $x$  annihilates a quark inside the proton. This quark carries a fraction  $\xi$  of the total proton momentum.

Similarly:

$$\frac{1}{2} \sum_{\text{spin}} \langle P(p) | -A_{\perp}^{\mu, R}(s\bar{u}) A_{\perp, \mu}^a(0) | P(p) \rangle$$

$$= \int_{-\frac{1}{2}}^{\frac{1}{2}} d\zeta f_{g/p}(\zeta) e^{i\zeta s\bar{u} \cdot p}$$

↙ gluon PDF
↘ charge-conjugation

$$f_{g/p}(-\zeta) = f_{g/p}(\zeta)$$

Since the soft fields decouple and  $L_c \cong L_{\text{qed}}$ , these SCET matrix elements are equivalent to the PDFs defined in QCD.

Plugging into  $T_{\mu\nu}$ , we get

$$T_{\mu\nu} = i \int d^4z e^{iqz} \dots$$




$$\begin{aligned}
& \langle P(p) | T \{ J_\mu(z) J_\nu(0) \} | P(p) \rangle \\
&= \int_0^1 d\xi \sum_q \tilde{C}_q^{\mu\nu}(\varphi^2, \bar{n} \cdot p \xi) \bar{n} \cdot p f_{q/p}(\xi) \\
&\quad + \text{"glue"}
\end{aligned}$$

To obtain  $\tilde{C}_q^{\mu\nu}$ , one performs a matching computation. The easiest way to obtain the coefficient is to work with partonic states, i.e. external quarks or gluons.

For an external quark of flavor  $q$  and momentum  $\xi \cdot p$ , we have

$$f_{q'/q}(\xi) = \delta(1-\xi) \delta_{qq'}$$

and the Wilson coefficient is equivalent to the diagram



In summary, one finds the following:


$$\sigma_{e^- p \rightarrow e^- X} = \sum_{i=q,\bar{q},g} \int_0^1 d\xi f_{i/p}(\xi) \cdot \hat{\sigma}_{e^- i \rightarrow e^- X}$$

partonic amplitude  
the incoming parton  
has momentum

$$\xi \cdot p$$

↑  
parton momentum

A similar result holds for  $pp$  collisions, but with two PDFs for two incoming partons, e.g.



Finally, let us remark that the PDF matrix elements suffer from UV divergences and need renormalization. The renormalized PDFs depend on the scale  $\mu$  and fulfill a RG equation

$$\frac{d}{d \ln \mu} f_{i/p}(\xi, \mu) = \sum_j P_{i \leftarrow j} \otimes f_j$$

$$= \sum_j \int_{\xi}^1 dx P_{i \leftarrow j} \left( \frac{\xi}{x}, \alpha_s(\mu) \right) f_{j/p}(x, \mu)$$

The different PDFs mix under renormalization.

The PDFs contain non-perturbative bound-state dynamics. There are methods to extract information on them using lattice QCD, but as of now most information on them is obtained by assuming a functional form (at some reference scale  $\mu_0$ ) and then fitting to experimentally measured cross sections.